

QED Radiation in Vincia

Rob Verheyen

With Peter Skands & Ronald Kleiss

Work in progress



Radboud University



Introduction

Vincia is a parton shower plugin for Pythia based on antenna factorization

Currently Vincia only does QCD radiation

We want to include QED radiation too

[Giele, Kosower, Skands:1102.2126](#)

[Gehrmann, Ritzmann, Skands:1108.6172](#)

Current approaches to photon radiation

DGLAP

- Resums collinear photon logarithms
- Interleaving with QCD shower

YFS

- Resums soft photon logarithms
- Collinear logarithms can be included, but not resummed
- Afterburner to add soft photons


I'll discuss the three algorithms for photon emission we're implementing

Leading Color Gluon Emission

Factorization

$$|M(\dots, p_a, k, \dots)|^2 \xrightarrow{p_a \parallel k} g^2 C \frac{P(z)}{p_a \cdot k} |M(\dots, p_a + k, \dots)|^2$$

$$|M(\dots, p_a, k, p_b, \dots)|^2 \xrightarrow{k \rightarrow 0} g^2 C \left[\frac{2p_a \cdot p_b}{(p_a \cdot k)(k \cdot p_b)} - \frac{m_a^2}{(p_a \cdot k)^2} - \frac{m_b^2}{(p_b \cdot k)^2} \right] |M(\dots, p_a, p_b, \dots)|^2$$




Leading Color Gluon Emission

Factorization

$$|M(\dots, p_a, k, p_b, \dots)|^2 \approx g^2 C a_e^{QCD}(p_a, k, p_b) |M(\dots, p'_a, p'_b, \dots)|^2$$

Ordering scale

$$t = p_{\perp}^2 = 4 \frac{p_a \cdot k p_b \cdot k}{m^2}$$



2 → 3 branching


Illustration: G. Salam

Photon Emission

Factorization

$$|M(\dots, p_a, k, \dots)|^2 \xrightarrow{p_a \parallel k} e^2 Q_a^2 \frac{P(z)}{p_a \cdot k} |M(\dots, p_a + k, \dots)|^2$$

$$|M(\{p\}, k)|^2 \xrightarrow{k \rightarrow 0} -e^2 \sum_{[a,b]} Q_a Q_b \left[\frac{2p_a \cdot p_b}{(p_a \cdot k)(k \cdot p_b)} - \frac{m_a^2}{(p_a \cdot k)^2} - \frac{m_b^2}{(p_b \cdot k)^2} \right] |M(\{p\})|^2$$



Photon Emission

Factorization

$$|M(\{p\}, k)|^2 \approx e^2 a_e^{QED}(\{p\}, k) |M(\{p'\})|^2$$

$$a_e^{QED}(\{p\}, k) = - \sum_{[a,b]} Q_a Q_b a_e^{QCD}(p_a, k, p_b)$$

$n \rightarrow n + 1$ branching

Ordering scale

$$t = p_{\perp}^2 = 4 \frac{p_a \cdot k p_b \cdot k}{m^2}$$

Photon emissions are a multi-scale problem

Goal: recast this $n \rightarrow n + 1$ branching into a (set of) $2 \rightarrow 3$ branchings

Option 1: Pairing

Incoherent Pairing

Pythia-like approach: Include only one antenna function for every fermion

$$a_e^{QED}(\{p\}, k) = Q_{f_1^+} Q_{f_1^-} a_e^{QCD}(p_{f_1^+}, k, p_{f_1^-}) + Q_{f_2^+} Q_{f_2^-} a_e^{QCD}(p_{f_2^+}, k, p_{f_2^-}) + \dots$$

Competition between independent radiators


- Correct collinear behaviour
- Only includes some eikonal factors

Pair up the fermions to minimise $m_{f_1^+ f_1^-}^2 + m_{f_2^+ f_2^-}^2 + \dots$

Incoherent Pairing

Photon radiation should decrease as the angle between opposite charges decreases

Emission scales are kinematically restricted by the antenna mass



Pair up the fermions to minimise $m_{f_1^+ f_1^-}^2 + m_{f_2^+ f_2^-}^2 + \dots$
Brute force $\mathcal{O}(n!)$ complexity...

The Hungarian Algorithm

Turns out this is a well-known problem from graph theory!

$$\begin{bmatrix} f_1^+ \\ f_2^+ \\ f_3^+ \end{bmatrix} \begin{bmatrix} f_1^- & f_2^- & f_3^- \\ m_{f_1^+ f_1^-}^2 & m_{f_1^+ f_2^-}^2 & m_{f_1^+ f_3^-}^2 \\ m_{f_2^+ f_1^-}^2 & m_{f_2^+ f_2^-}^2 & m_{f_2^+ f_3^-}^2 \\ m_{f_3^+ f_1^-}^2 & m_{f_3^+ f_2^-}^2 & m_{f_3^+ f_3^-}^2 \end{bmatrix}$$

The Hungarian Algorithm

Let's look at an example to see how it works

$$\begin{array}{c} \left[\begin{array}{c} f_1^+ \\ f_2^+ \\ f_3^+ \end{array} \right] \end{array} \begin{array}{c} \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \end{array} \right] \\ \left[\begin{array}{ccc} 35 & 50 & 30 \\ 5 & 15 & 10 \\ 35 & 50 & 20 \end{array} \right] \end{array}$$

The Hungarian Algorithm

Step 1: Subtract the lowest row element from all rows

$$\begin{array}{c} [f_1^- \quad f_2^- \quad f_3^-] \\ [f_1^+ \quad f_2^+ \quad f_3^+] \end{array} \begin{bmatrix} 5 & 20 & 0 \\ 0 & 10 & 5 \\ 15 & 30 & 0 \end{bmatrix} \begin{array}{l} -30 \\ -5 \\ -20 \end{array}$$

The Hungarian Algorithm

Step 2: Subtract the lowest column element from all rows

$$\begin{array}{c} \left[\begin{array}{c} f_1^+ \\ f_2^+ \\ f_3^+ \end{array} \right] \end{array} \quad \begin{array}{c} \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \end{array} \right] \\ \left[\begin{array}{ccc} 5 & 10 & 0 \\ 0 & 0 & 5 \\ 15 & 20 & 0 \end{array} \right] \\ -10 \end{array}$$

The Hungarian Algorithm

Step 3: Find the minimal line covering

$$\begin{array}{c} \left[\begin{array}{c} f_1^+ \\ f_2^+ \\ f_3^+ \end{array} \right] \end{array} \begin{array}{c} \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \\ \hline 5 & 10 & 0 \\ 0 & 0 & 5 \\ \hline 15 & 20 & 0 \end{array} \right] \end{array}$$

If the line covering is maximal ($n=3$), pairing with cost 0 can be found

The Hungarian Algorithm

Step 4: Find the lowest uncovered element

$$\begin{array}{c} \left[\begin{array}{c} f_1^+ \\ f_2^+ \\ f_3^+ \end{array} \right] \end{array} \begin{array}{c} \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \\ \hline 5 & 10 & 0 \\ 0 & 0 & 5 \\ 15 & 20 & 0 \end{array} \right] \end{array}$$

The Hungarian Algorithm

Step 4: Subtract that number from all uncovered element
Add it to all doubly covered elements

$$\begin{array}{c} -5 \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \end{array} \right] \\ \left[\begin{array}{c} f_1^+ \\ f_2^+ \\ f_3^+ \end{array} \right] \left[\begin{array}{ccc} 0 & 5 & 0 \\ 0 & 0 & 10 \\ 10 & 15 & 0 \end{array} \right] +5 \end{array}$$

And go back to step 3

The Hungarian Algorithm

$$\begin{array}{c} \left[\begin{array}{ccc} f_1^- & f_2^- & f_3^- \end{array} \right] \\ \left[\begin{array}{ccc} f_1^+ & f_2^+ & f_3^+ \end{array} \right] \end{array} \begin{array}{c} \left[\begin{array}{ccc} \textcircled{0} & 5 & 0 \\ 0 & \textcircled{0} & 10 \\ 10 & 15 & \textcircled{0} \end{array} \right] \end{array}$$

Now we are able to find an optimal pairing!

$\mathcal{O}(n^3)$ complexity, so not computationally prohibitive

Option 2: Coherent

Coherent Emission

Separate phase space into sectors

$$|M(\{p\}, k)|^2 \approx \sum_{[i,j]} \left(- \sum_{[a,b]} Q_a Q_b a_e^{QCD}(p_a, k, p_b) \right) \theta(p_{\perp ij}^2) |M(\dots, p'_i, p'_j, \dots)|^2$$

2 → 3 branching



1 if $p_{\perp ij}^2$ is the smallest
0 otherwise



Equivalent to ordering in

$$t = \min(p_{\perp ij}^2) = \min\left(4 \frac{p_i \cdot k p_j \cdot k}{m^2}\right)$$

But there's a problem...

Sudakov Veto Algorithm

Want to sample from

$$f(t) \exp\left(-\int_t^u d\tau f(\tau)\right)$$

Find $g(t) \geq f(t)$

Set $u = t_{\text{start}}$

Sample t from $g(t) \exp\left(-\int_t^u d\tau g(\tau)\right)$

Set $u = t$

Accept with probability $\frac{f(t)}{g(t)}$



Done $\rightarrow t$

Coherent Emission

We need an overestimate for the branching kernel

$$a_e^{QED} = - \sum_{[a,b]} Q_a Q_b a_e^{QCD}(p_a, k, p_b)$$

It's possible to find one, but...



The algorithm is slow!

Option 3: Coherent Weighted

Sudakov Veto Algorithm

Want to sample from

$$f(t) \exp\left(-\int_t^u d\tau f(\tau)\right)$$

Find $g(t) \not\geq f(t)$

Set $u = t_{\text{start}}$

Sample t from $g(t) \exp\left(-\int_t^u d\tau g(\tau)\right)$

Set $u = t$

Accept with probability $p(t)$

Apply weight

$$\frac{1}{p(t)} \frac{f(t)}{g(t)}$$

Done $\rightarrow t$



Apply weight

$$\frac{1}{g(t)} \frac{g(t) - f(t)}{1 - p(t)}$$

Coherent Weighted Emission

Use an event-based incomplete overestimate

$$p(t) = \tanh \left(\frac{f(t)}{g(t)} \right)$$



$$R = - \sum_{[a,b]} Q_a Q_b (1 - \cos(\theta_{ab}))$$

Much faster, but events are weighted

Summary

We're implementing three ways of doing photon emissions

1. Incoherent Pairing

- Fast
- Not coherent, but has most important eikonals

2. Coherent Unweighted

- Slow
- Fully coherent



3. Coherent Weighted

- Fast
- Weighted events

Extra Slides



Comparison - Coherence



Comparison - DGLAP equation

